

Second order SUSY transformations with ‘complex energies’

David J. Fernández C.^{a,1}, Rodrigo Muñoz^a and
Arturo Ramos^{a,b,2}

^a*Departamento de Física, CINVESTAV, A.P. 14-740, 07000 México D.F., Mexico*

^b*Dipartimento di Matematica Pura ed Applicata, Università degli Studi di Padova,
Via G. Belzoni 7, I-35131 Padova, Italy*

Abstract

Second order supersymmetry transformations which involve a pair of complex conjugate factorization energies and lead to real non-singular potentials are analyzed. The generation of complex potentials with real spectra is also studied. The theory is applied to the free particle, one-soliton well and one-dimensional harmonic oscillator.

Key words: Second-order supersymmetry, irreducible intertwining operators, complex potentials with real spectra, generation of solvable potentials

PACS: 03.65.Ca, 03.65.Fd, 03.65.Ge, 11.30.Pb

1 Introduction

The n -th order supersymmetric quantum mechanics (n -SUSY QM), which involves differential intertwining operators of order n , is a useful tool for generating new solvable potentials [1, 2, 3, 4]. Due to its simplicity, the 1-SUSY QM is the most explored; its nonsingular transformations produce partner potentials whose spectra can differ at most in the ground state energy level [5, 6, 7]. The

Email addresses: david@fis.cinvestav.mx (David J. Fernández C.), rodrigom@fis.cinvestav.mx (Rodrigo Muñoz), aramos@math.unipd.it (Arturo Ramos).

¹ Corresponding author. Tf: +52 57473879. Fax: +52 57473879.

² Work developed in part while A.R. was visiting the Dept. de Física of CINVESTAV in the fall of 2000. On leave of absence from Departamento de Física Teórica, Facultad de Ciencias, Universidad de Zaragoza, 50009 Zaragoza, Spain.

difficulty of ‘modifying’ the excited part of the spectrum has been surpassed through the 2-SUSY QM [8,9,10,11,12,13,14], which allows to ‘create’ two new levels ϵ_1, ϵ_2 between two neighboring energies E_i, E_{i+1} of the initial Hamiltonian [15]. A similar treatment, implemented for periodic potentials [16], can be used to embed two bound states in a spectral gap above the lowest energy band [17, 18]. In both situations the corresponding 2-SUSY transformations are *irreducible*, i.e., when obtained as the iteration of two 1-SUSY procedures they will involve always singular intermediate potentials.

Here we will study a different set of ‘irreducible’ 2-SUSY transformations applied to non-periodic potentials, which employs two complex conjugate factorization energies $\epsilon_1, \epsilon_2, \epsilon_2 = \bar{\epsilon}_1$. Irreducibility means now that the intermediate potential is complex although the final one is real. This problem has been addressed previously [19], but up to our knowledge the conditions granting that the final potential will be regular have not been yet examined. We will show as well that the non-singular case leads to intermediate complex potentials having real spectra. These points constitute the subject of this letter, which has been organized as follows. In section 2 the second order SUSY transformations with $\epsilon_1, \epsilon_2 \in \mathbb{C}, \epsilon_2 = \bar{\epsilon}_1$ will be analyzed. A prescription for avoiding the singularities in the new potential will be provided in section 3, while section 4 will be devoted to study the intermediate complex potentials. Section 5 will deal with some particular examples as the free particle, one-soliton well and harmonic oscillator potential.

2 Second order supersymmetric quantum mechanics

The standard supersymmetric (SUSY) algebra with generators Q_1, Q_2 (supercharges) and H_{ss} (SUSY Hamiltonian) reads:

$$\{Q_j, Q_k\} = \delta_{jk} H_{ss}, \quad [H_{ss}, Q_j] = 0, \quad j, k = 1, 2, \quad (1)$$

where $[\cdot, \cdot]$ denotes a commutator and $\{\cdot, \cdot\}$ an anticommutator. The realization $Q_1 = (Q^\dagger + Q)/\sqrt{2}, Q_2 = (Q^\dagger - Q)/(i\sqrt{2})$ with

$$Q = \begin{pmatrix} 0 & A \\ 0 & 0 \end{pmatrix}, \quad Q^\dagger = \begin{pmatrix} 0 & 0 \\ A^\dagger & 0 \end{pmatrix}, \quad (2)$$

$$H_{ss} = \begin{pmatrix} AA^\dagger & 0 \\ 0 & A^\dagger A \end{pmatrix} = \begin{pmatrix} (\widetilde{H} - \epsilon_1)(\widetilde{H} - \epsilon_2) & 0 \\ 0 & (H - \epsilon_1)(H - \epsilon_2) \end{pmatrix}, \quad (3)$$

$$A = \frac{d^2}{dx^2} + \eta(x) \frac{d}{dx} + \gamma(x), \quad (4)$$

is called second order supersymmetric quantum mechanics (2-SUSY QM). In this formalism H , $\tilde{H}$ are two intertwined Schrödinger Hamiltonians:

$$\tilde{H}A = AH, \quad (5)$$

$$H = -\frac{d^2}{dx^2} + V(x), \quad \tilde{H} = -\frac{d^2}{dx^2} + \tilde{V}(x), \quad (6)$$

and thus the real functions $\eta(x)$, $\gamma(x)$ are related with $V(x)$, $\tilde{V}(x)$ through:

$$\tilde{V} - V = 2\eta', \quad (7)$$

$$(\tilde{V} - V)\eta = 2V' + 2\gamma' + \eta'', \quad (8)$$

$$(\tilde{V} - V)\gamma = \eta V' + V'' + \gamma''. \quad (9)$$

To decouple this system, substitute (7) in (8) and integrate to obtain

$$\gamma = d - V + \eta^2/2 - \eta'/2, \quad (10)$$

where $d \in \mathbb{R}$ is a constant. By plugging (7,10) into (9), multiplying the result by η and performing the next integration one arrives to:

$$\eta\eta'' - (\eta')^2/2 + 2\eta^2\left(\eta^2/4 - \eta' - V + d\right) + 2c = 0, \quad (11)$$

$c \in \mathbb{R}$ being another constant. This formalism is useful for generating new solvable potentials. To illustrate it, suppose that c, d are fixed and $V(x)$ is an initial exactly solvable potential. The complete determination of $\tilde{V}(x)$ in (7) requires to find solutions $\eta(x)$ of the nonlinear second order differential equation (11). Let us find them using the *Ansatz* [12]:

$$\eta'(x) = \eta^2(x) + 2\beta(x)\eta(x) - 2\xi(x), \quad (12)$$

where $\beta(x)$ and $\xi(x)$ are to be determined. After substituting (12) into (11) we obtain a system of equations from which it follows that $\xi^2 = c$. The essential part of the system is the Riccati equation:

$$\beta' + \beta^2 = V - \epsilon, \quad \epsilon = d + \xi. \quad (13)$$

As there exist two possible values of ϵ , $\epsilon_1 = d + \sqrt{c}$ and $\epsilon_2 = d - \sqrt{c}$ (which coincide with the factorization energies in (3)), we indeed are dealing with two equations (13) whose solutions will be denoted by $\beta_1(x)$, $\beta_2(x)$. This leads to a natural classification of the 2-SUSY transformations based on the sign of c ,

which will be discussed elsewhere. Here, we restrict ourselves to the *complex* case for which $c < 0$ and then $\epsilon_1, \epsilon_2 \in \mathbb{C}$, $\epsilon_2 = \bar{\epsilon}_1$. Since V is real we can take $\beta_2(x) = \bar{\beta}_1(x)$, i.e., the problem reduces to solve the Riccati equation for β_1 . In addition, using (12) we get two equivalent expressions for the real $\eta(x)$:

$$\eta' = \eta^2 + 2\beta_1\eta - 2i\text{Im}(\epsilon_1), \quad (14)$$

$$\eta' = \eta^2 + 2\bar{\beta}_1\eta + 2i\text{Im}(\epsilon_1). \quad (15)$$

By subtracting both equations and solving for η , we obtain:

$$\eta = \text{Im}(\epsilon_1)/\text{Im}(\beta_1). \quad (16)$$

Once we know η , the 2-SUSY partner potential $\tilde{V}(x)$ is calculated using

$$\tilde{V} = V + 2 [\text{Im}(\epsilon_1)/\text{Im}(\beta_1)]'. \quad (17)$$

Let us remark that the case we are dealing with has been previously explored [8]. However, we have not found any previous analysis on how to avoid the singularities in $\tilde{V}(x)$, a phenomenon which seems almost unavoidable in the complex case [19].

3 The non-singular 2-SUSY potentials

The simplest algorithms departing from and arriving at the exactly solvable potentials should avoid the singularities which might appear in $\tilde{V}(x)$. Notice that a singular $\tilde{V}(x)$ could be treated as a non-singular partner potential of $V(x)$ in a restricted x -domain. However, this would require at the end to solve the initial Schrödinger equation with modified boundary conditions loosing, in general, the solvability of H [20].

Let us rewrite first the formulae of section 2 in terms of solutions $u_1(x)$ of the Schrödinger equation arising from (13) by the change $\beta(x) = u'(x)/u(x)$ [14]:

$$-u''(x) + V(x)u(x) = \epsilon u(x). \quad (18)$$

Hence $\eta(x) = -2i\text{Im}(\epsilon_1)|u_1|^2/W(u_1, \bar{u}_1)$, where $W(u_1, u_2) = u_1u_2' - u_2u_1'$ denotes the Wronskian of u_1, u_2 . From now on it is convenient to work with the real *normalized Wronskian* $w(x) \equiv W(u_1, \bar{u}_1)/[2i\text{Im}(\epsilon_1)]$. Therefore:

$$w'(x) = |u_1(x)|^2, \quad (19)$$

$$\eta(x) = -w'(x)/w(x), \quad (20)$$

$$\tilde{V}(x) = V(x) - 2[w'(x)/w(x)]'. \quad (21)$$

In order that $\tilde{V}(x)$ be non-singular, $w(x)$ must be nodeless. Since $w(x)$ is an increasing monotonic function (see (19)), the arising of zeros is avoided if

$$\lim_{x \rightarrow \infty} w(x) = 0 \quad \text{or} \quad \lim_{x \rightarrow -\infty} w(x) = 0. \quad (22)$$

To ensure this requirement it is sufficient that either

$$\lim_{x \rightarrow \infty} u_1(x) = 0 \quad \text{or} \quad \lim_{x \rightarrow -\infty} u_1(x) = 0. \quad (23)$$

Such solutions are appropriate for generating non-singular potentials $\tilde{V}(x)$. Notice that a similar treatment can be designed for systems defined in a generic interval $x \in (a, b) \subset \mathbb{R}$ by identifying in (22-23) $-\infty$ with a and ∞ with b .

4 Complex potentials with real spectrum

Although in principle irreducible, let us decompose the non-singular 2-SUSY transformations of the previous section into two 1-SUSY steps:

$$\tilde{H}A_2 = A_2H_1, \quad H_1A_1 = A_1H, \quad (24)$$

where

$$H_1 = -\frac{d^2}{dx^2} + V_1(x), \quad A_i = \frac{d}{dx} + \alpha_i(x), \quad i = 1, 2. \quad (25)$$

The 1-SUSY treatment implies that α_1, α_2 obey the Riccati equations:

$$-\alpha_1' + \alpha_1^2 = V(x) - \epsilon_1, \quad (26)$$

$$-\alpha_2' + \alpha_2^2 = V_1(x) - \bar{\epsilon}_1, \quad (27)$$

where $V_1(x) = V(x) + 2\alpha_1'$ and thus $\tilde{V}(x) = V(x) + 2(\alpha_1 + \alpha_2)'$. A simple comparison of (13) with (26) leads to

$$\alpha_1(x) = -\beta_1(x) = -u_1'(x)/u_1(x), \quad (28)$$

$u_1(x)$ being a solution of (18) behaving asymptotically as in (23). Moreover, by expanding $A = A_2 A_1$ and comparing the result with (4) we find that:

$$\alpha_2 = -\alpha_1 + \eta = \beta_1 + (\epsilon_1 - \bar{\epsilon}_1)/(\beta_1 - \bar{\beta}_1). \quad (29)$$

This is a particular case of the finite difference Bäcklund algorithm [3, 13, 14], which algebraically determines a solution to (27) in terms of solutions of (26) for two different factorization energies (here ϵ_1 and $\bar{\epsilon}_1$). It is interesting as well to factorize the involved Hamiltonians:

$$H = A_1^- A_1 + \epsilon_1, \quad H_1 = A_1 A_1^- + \epsilon_1, \quad (30)$$

$$H_1 = A_2^- A_2 + \bar{\epsilon}_1, \quad \widetilde{H} = A_2 A_2^- + \bar{\epsilon}_1, \quad (31)$$

$A_i^- = -d/dx + \alpha_i(x)$, $i = 1, 2$. Since α_1, α_2 and ϵ_1 are complex, the A_i^- 's are not adjoint to the A_i 's but $A_i^\dagger = -d/dx + \bar{\alpha}_i(x)$, $i = 1, 2$.

It is clear now that the complex intermediate potential $V_1(x)$ is non-singular:

$$V_1(x) = V(x) - 2[u_1'(x)/u_1(x)]'. \quad (32)$$

To analyse the normalizability of the corresponding eigenfunction associated to E_n ,

$$\psi_n^1(x) = c_n A_1 \psi_n = c_n [\psi_n'(x) - u_1'(x) \psi_n(x)/u_1(x)], \quad (33)$$

we will employ the operator relationship:

$$\eta A_1 = H - \epsilon_1 + A. \quad (34)$$

From the validity of (23) and the assumption of $||\psi_n|| = 1$, it turns out that $\tilde{\psi}_n = A\psi_n/|E_n - \epsilon_1|$ is normalized, and therefore the function $\eta A_1 \psi_n = (E_n - \epsilon_1)\psi_n + |E_n - \epsilon_1|\tilde{\psi}_n$ is normalizable as well. Thus, for $A_1 \psi_n$ to be normalizable it is necessary that η^{-1} does not destroy the normalizability of $(E_n - \epsilon_1)\psi_n + |E_n - \epsilon_1|\tilde{\psi}_n$. If this is the case (and this will happen for the examples we discuss below), we obtain a complex potential $V_1(x)$ with real eigenvalues E_n [21, 22, 23]. Let us remark that complex Hamiltonians with real spectra have been studied recently in the context of PT-symmetry and pseudo-Hermiticity [24, 25].

5 Illustrative examples

We shall show that the previous techniques admit very simple applications.

i) Consider firstly the free particle for which $V(x) = 0$. The general solution $u_1(x)$ of the Schrödinger equation (18) for $\epsilon_1 \in \mathbb{C}$ is a linear combination of

$$e^{\pm(k_1+ik_2)x}, \quad (35)$$

where $\epsilon_1 = -(k_1+ik_2)^2$, $k_1 > 0$, $k_2 \in \mathbb{R}$. In general, such a $u_1(x)$ does not tend to zero neither when $x \rightarrow -\infty$ nor when $x \rightarrow +\infty$. However, two particular solutions with the required behavior are precisely those of (35). We use them for obtaining the nodeless $w(x)$:

$$w(x) = \pm e^{\pm 2k_1 x} / (2k_1). \quad (36)$$

It turns out that $\tilde{V}(x)$ becomes again the null potential for both $w(x)$, $\tilde{V}(x) = 0$. The intermediate 1-SUSY complex potentials generated by using the two $u_1(x)$ of (35) are as well trivial, $V_1(x) = 0$. Our conclusion is that the null potential can be non-trivially transformed in frames of our algorithm only at the price of creating singularities (compare with [19]).

ii) Consider now the well known one-soliton potential (Pöschl-Teller) [26]

$$V(x) = -2k_0^2 \text{sech}^2(k_0 x) \quad (37)$$

which is obtained from the null potential by a 1-SUSY transformation employing $\cosh(k_0 x)$, $k_0 > 0$. The spectrum of (37) consists of a continuous part $E \geq 0$ and a bound state at $E_0 = -k_0^2$ with eigenfunction given by:

$$\psi_0(x) = \sqrt{k_0/2} \text{sech}(k_0 x). \quad (38)$$

The general solution $u_1(x)$ of (18) for (37) with $\epsilon_1 = -(k_1+ik_2)^2$, $k_1 > 0$, $k_2 \in \mathbb{R}$ is a linear combination of the 1-SUSY transformed eigenfunctions of (35)

$$e^{\pm(k_1+ik_2)x} [k_0 \tanh(k_0 x) \mp (k_1 + ik_2)]. \quad (39)$$

The solutions (39) are precisely the required ones: if the upper signs are taken, then $u_1(x) \rightarrow 0$ for $x \rightarrow -\infty$, while the lower signs ensure $u_1(x) \rightarrow 0$ when $x \rightarrow +\infty$. An explicit calculation leads to the two nodeless $w(x)$:

$$w(x) = \pm \frac{ke^{\pm 2k_1 x} \cosh[k_0(x \mp x_0)]}{2k_1 \cosh(k_0 x)}, \quad (40)$$

where $k_0^2 + k_1^2 + k_2^2 \equiv k \cosh(k_0 x_0)$, $2k_0 k_1 \equiv k \sinh(k_0 x_0)$, $k, x_0 \in \mathbb{R}$. The two 2-SUSY partner potentials of (37) read now:

$$\tilde{V}(x) = -2k_0^2 \text{sech}^2[k_0(x \mp x_0)], \quad (41)$$

obtaining just real x_0 -displaced copies of (37). This result has to do with the Darboux invariance phenomenon recently discovered for the one-soliton well [17, 18].

On the other hand, the two complex intermediate potentials generated by (39) become:

$$V_1(x) = -2k_0^2 \text{sech}^2[k_0(x \mp x_1)], \quad (42)$$

where now $k_1 + ik_2 \equiv \kappa \cosh(k_0 x_1)$, $k_0 \equiv \kappa \sinh(k_0 x_1)$ define ‘complex displacements’ $x_1 \in \mathbb{C}$, $\kappa \in \mathbb{C}$. These potentials have a bound state at $E_0 = -k_0^2$ whose normalized ‘ground state’ eigenfunction is obtained from (33) by employing the $\psi_0(x)$ of (38) and the $u_1(x)$ of (39):

$$\psi_0^1(x) = \frac{k_0}{|\kappa|} \left[\frac{1}{k_2} \arctan\left(\frac{k_0 + k_1}{k_2}\right) + \frac{1}{k_2} \arctan\left(\frac{k_0 - k_1}{k_2}\right) \right]^{-\frac{1}{2}} \text{sech}[k_0(x \mp x_1)] \quad (43)$$

The complex potentials (42) were obtained for the first time in [22].

iii) Our final example is the harmonic oscillator:

$$V(x) = x^2, \quad (44)$$

which has a purely discrete equidistant spectrum $\{E_n = 2n + 1, n = 0, 1, \dots\}$. The general solution of (18) for $\epsilon_1 \in \mathbb{C}$ is now (see [4] and references therein):

$$u_1(x) = c_1 e^{-\frac{x^2}{2}} \left[{}_1F_1\left(\frac{1 - \epsilon_1}{4}, \frac{1}{2}; x^2\right) + 2\nu x \frac{\Gamma(\frac{3 - \epsilon_1}{4})}{\Gamma(\frac{1 - \epsilon_1}{4})} {}_1F_1\left(\frac{3 - \epsilon_1}{4}, \frac{3}{2}; x^2\right) \right], \quad (45)$$

where ${}_1F_1(a, c; z)$ is the Kummer hypergeometric series. In general, such a $u_1(x)$ does not satisfy neither $\lim_{x \rightarrow -\infty} u_1(x) = 0$ nor $\lim_{x \rightarrow \infty} u_1(x) = 0$. However, there are two particular values for ν ($\nu = \pm 1$) leading to solutions with the required behavior:

$$u_1(x) = e^{-\frac{x^2}{2}} \left[{}_1F_1\left(\frac{1-\epsilon_1}{4}, \frac{1}{2}; x^2\right) \pm 2x \frac{\Gamma(\frac{3-\epsilon_1}{4})}{\Gamma(\frac{1-\epsilon_1}{4})} {}_1F_1\left(\frac{3-\epsilon_1}{4}, \frac{3}{2}; x^2\right) \right]. \quad (46)$$

Take, e.g., (46) with the upper sign, which in the negative semiaxis $x = -|x| < 0$ reduces to:

$$u_1(x) = \frac{\Gamma(\frac{3-\epsilon_1}{4})}{\Gamma(\frac{1}{2})} e^{-\frac{|x|^2}{2}} \Psi\left(\frac{1-\epsilon_1}{4}, \frac{1}{2}; |x|^2\right), \quad (47)$$

where the Tricomi function $\Psi(a, c; z)$ is related with ${}_1F_1(a, c; z)$ through (see, e.g., [27]):

$$\Psi(a, c; z) = \frac{\Gamma(1-c)}{\Gamma(a-c+1)} {}_1F_1(a, c; z) + \frac{\Gamma(c-1)}{\Gamma(a)} z^{1-c} {}_1F_1(a-c+1, 2-c; z) \quad (48)$$

Since the leading term in the asymptotic expansion for $\Psi(a, c; z)$ is z^{-a} [28], one can check that $\lim_{x \rightarrow -\infty} u_1(x) = 0$. Similarly, if the lower sign is chosen we have $\lim_{x \rightarrow \infty} u_1(x) = 0$.




Fig. 1. The real potential $\tilde{V}(x)$ (black curve) generated from $V(x) = x^2$ (gray curve) by means of a ‘complex’ 2-SUSY transformation with $\epsilon_1 = 10 + 0.1i$ and the $u_1(x)$ of (46) with the lower minus sign.

Once we have identified the solutions (46) with the right asymptotic behavior, we evaluated $w(x)$ and then $\tilde{V}(x)$. The resulting expressions in this case are too long; instead, we are plotting $\tilde{V}(x)$ for $\epsilon_1 = 10 + 0.1i$ using (46) with the lower minus sign (see figure 1). Contrasting with the results for the previous examples, in this case the potentials $\tilde{V}(x)$ are in general different from $V(x)$. This means that the transformations involving a pair of complex conjugate factorization energies are effective tools in generating isospectral 2-SUSY partner potentials. As a byproduct, we have obtained in a simple way complex potentials $V_1(x)$ given by (32) with real energy eigenvalues $E_n = 2n + 1$. A plot of the ‘ground state’ probability density $|\psi_0^1(x)|^2$, illustrating the existence of these bound states for the complex 1-SUSY partner $V_1(x)$ of the oscillator, is shown in figure 2.




Fig. 2. The ‘ground state’ probability density $|\psi_0^1(x)|^2$ for the complex 1-SUSY partner potential (32) of the oscillator generated by using $u_1(x)$ of (46) with the lower sign and $\epsilon_1 = 10 + 0.1i$.

6 Conclusions

We have shown that the 2-SUSY transformations involving two complex conjugate factorization energies can produce new non-singular potentials isospectral to a given initial one. This non-singular character is shared as well by the intermediate complex potentials arising when those transformations are factorized.

Acknowledgements. The support of CONACYT (México) is acknowledged. A.R. has been partially supported by the Spanish Ministerio de Educación y Cultura through a FPI grant, research project PB96-0717, and by the European Commission funding for the Research Training Network “Mechanics and Symmetry in Europe” (MASIE), contract HPRN-CT-2000-00113. A.R. thanks also the warm hospitality at the Departamento de Física of CINVESTAV.

References

- [1] A.A. Andrianov, M.V. Ioffe, V.P. Spiridonov, Phys. Lett. A **174** (1993) 273.
- [2] V.G. Bagrov, B.F. Samsonov, Phys. Part. Nucl. **28** (1997) 374.
- [3] D.J. Fernández, V. Hussin, B. Mielnik, Phys. Lett. A **244** (1998) 309.
- [4] D.J. Fernández, V. Hussin, J. Phys. A **32** (1999) 3603.
- [5] J.F. Cariñena, G. Marmo, A.M. Perelomov, M.F. Rañada, Int. J. Mod. Phys. A **13** (1998) 4913.
- [6] J.F. Cariñena, A. Ramos, Mod. Phys. Lett. A **15** (2000) 1079.
- [7] J.F. Cariñena, A. Ramos, Rev. Math. Phys. **12** (2000) 1279.

- [8] A.A. Andrianov, M.V. Ioffe, F. Cannata, J.P. Dedonder, Int. J. Mod. Phys. A **10** (1995) 2683.
- [9] A.A. Andrianov, M.V. Ioffe, D.N. Nishnianidze, Phys. Lett. A **201** (1995) 103.
- [10] D.J. Fernández, Int. J. Mod. Phys. A **12** (1997) 171.
- [11] D.J. Fernández, M.L. Glasser, L.M. Nieto, Phys. Lett. A **240** (1998) 15.
- [12] O. Rosas-Ortiz, J. Phys. A **31** (1998) 10163; *ibid* **31** (1998) L507.
- [13] B. Mielnik, L.M. Nieto, O. Rosas-Ortiz, Phys. Lett. A. **269** (2000) 70.
- [14] J.F. Cariñena, A. Ramos, D.J. Fernández, Ann. Phys. (N.Y.) **292** (2001) 42.
- [15] B.F. Samsonov, Phys. Lett. A **263** (1999) 274.
- [16] D.J. Fernández, J. Negro, L.M. Nieto, Phys. Lett. A **275** (2000) 338.
- [17] D.J. Fernández, B. Mielnik, O. Rosas-Ortiz, B.F. Samsonov, Phys. Lett. A **294** (2002) 168.
- [18] D.J. Fernández, B. Mielnik, O. Rosas-Ortiz, B.F. Samsonov, J. Phys. A **35** (2002) 4279.
- [19] V.G. Bagrov, I.N. Ovcharov, B.F. Samsonov, J. Moscow Phys. Soc. **5** (1995) 191.
- [20] I.F. Márquez, J. Negro, L.M. Nieto, J. Phys. A **31** (1998) 4115.
- [21] F. Cannata, G. Junker, J. Trost, Phys. Lett. A **246** (1998) 219.
- [22] A.A. Andrianov, M.V. Ioffe, F. Cannata, J.P. Dedonder, Int. J. Mod. Phys. A **14** (1999) 2675.
- [23] B. Bagchi, S. Mallik, C. Quesne, Int. J. Mod. Phys. A **16** (2001) 2859; *ibid* **17** (2002) 51.
- [24] C.M. Bender, S. Boettcher, Phys. Rev. Lett. **80** (1998) 5243.
- [25] A. Mostafazadeh, J. Math. Phys. **43** (2002) 205, 2814, 3944.
- [26] J.I. Díaz, J. Negro, L.M. Nieto, O. Rosas-Ortiz, J. Phys. A **32** (1999) 8447.
- [27] P. Denner, A. Krzywicki, Mathematics for Physicists, Dover, New York, 1996.
- [28] M. Abramowitz, I.A. Stegun (Eds.), Handbook of Mathematical Functions, Wiley, New York, 1972.